

String theory T-duality and the zero point length of spacetime

A. Smailagic*, E. Spallucci†, T. Padmanabhan‡

*Sezione INFN di Trieste,
Strada Costiera 11, 34014 Trieste,
Italy**

*Department of Theoretical Physics,
University of Trieste, Strada Costiera 11, 34014 Trieste,
Italy, and*

*Sezione INFN di Trieste,
Strada Costiera 11, 34014 Trieste,
Italy† and*

IUCAA, Post Bag 4, Ganeshkhind, Pune - 411 007‡

(Dated: February 1, 2008)

It has been often conjectured that the correct theory of quantum gravity will act as a UV regulator in the low energy limit of quantum field theory. Earlier work has shown that if the path integral defining the quantum field theory propagator is modified, so that the amplitude is invariant under the duality transformation $l \rightarrow 1/l$ where l is the length of the path, then the propagator is UV-finite and exhibits a “zero-point length” of the spacetime. Since string theory uses extended structures and has a T-duality, these results should also emerge directly from string theory. We show, by explicit path integral computation, that this is indeed the case. The lowest order string theory correction to the propagator is the same as that obtained by the hypothesis of path integral duality.

Conventional quantum field theory exhibits UV divergences and it was always thought that this behavior will be “cured” in the correct theory of quantum gravity [1]. This is to be expected since one cannot operationally define scales below Planck scale, in any model that incorporates the principles of quantum theory and gravity [2], making Planck length act as the “zero-point length” of the spacetime. The propagator in a field theory should therefore be ultraviolet finite when quantum gravitational corrections are incorporated.

It was shown in [3] that if the path integral amplitude $\exp(-l(x, y))$ used for the definition of the Euclidean propagator is modified so that it is invariant under the duality transformation $l \rightarrow L_P^2/l$, then (i) the propagator becomes UV finite and (ii) the result is identical to that obtained by changing the interval $(x - y)^2$ to $(x - y)^2 + 4L_P^2$. This shows a deep connection between the duality $l \rightarrow L_P^2/l$ and the existence of zero-point length, which was not clearly understood.

The string theory has both these features (T-duality as well as finite length scale) built into it in a natural fashion. This suggests that one should be able to obtain similar results in standard string theory. In this letter we show that this is indeed the case: the zero-point length can be reproduced using T -duality of the string theory and the leading order corrections to the propagator, defined in a specific but consistent way, is the same as that obtained from the principle of path integral duality used in [3].

Since it is difficult (and unnecessary for our purpose) to obtain the low energy quantum field theory description directly from *full* string theory, we shall merely invoke certain key ingredients of the string theory to achieve our goal. There are three technical issues which are rel-

evant to establishing such a correspondence between the language of string theory and the low energy quantum field theory, so that one can sensibly ask for the corrections to the low energy propagator from string theory. The first point is that the result we want to derive is non-perturbative so that the simplest procedure, of doing this in a perturbative manner, is inadequate. The corrections to the propagator from zero-point length typically involves terms like $[x^2 + L_P^2]^{-1}$. This quantity is finite as $x \rightarrow 0$ but if we expand it in powers of L_P^2/x^2 , each term of the expansion will diverge as $x \rightarrow 0$. So we need to approach the problem non-perturbatively. Second, we need to identify a suitable (collective) degree of freedom of the string theory, which could be thought of as analogous to the low energy concept of a particle which is propagating. We find that the string center of mass (SCM) is a suitable candidate for this role and one can define a path integral propagator for the same. This propagator is UV finite because of the winding modes in the compact dimensions. Finally, we also need a mechanism to incorporate the light degrees of freedom (with masses far lower than the Planck mass) so that one can talk about a propagator for, say, an electron. This is possible by incorporating the effects of light modes by adding a suitable mass term in the Schwinger’s proper time formulation of the propagator. We shall now present the details of this approach.

Closed strings have the unique property to wrap around compactified spatial dimensions. In the case of a single compact dimension, the string mass spectrum is labeled, in addition to the harmonic excitation occupational number, by two more quantum numbers. The number n which determines the Kaluza-Klein excitation level and the number w which counts the string winding

around compactified dimensions. The string mass spectrum is given by [4]:

$$M^2 = \frac{1}{2\alpha'} \left(n^2 \frac{\alpha'}{R^2} + w^2 \frac{R^2}{\alpha'} \right) + \dots \quad (1)$$

where, $1/\alpha'$ and R are the string tension and the compactification radius respectively. (The dots refer to harmonic oscillators contributions which are irrelevant for our path integral calculation; some relevant features of the low energy excitations will be incorporated at the end of the calculation indirectly.) For the sake of simplicity, we have written (1) for the case of a single compact dimension (with the fifth-dimension compactified on a circle of length $l_0 = 2\pi R$) but our analysis can be extended to the case of D dimensional hyper-torus in a straightforward way. (We quote the result for the general case in the end). The important property of the spectrum (1) is its invariance under so-called T -duality. It is defined as a transformation relating the compactification radius and its inverse, while, at the same time, exchanging quantum numbers n and w as: $R \longleftrightarrow \alpha'/R$, $n \longleftrightarrow w$. This symmetry relates large and small distances, and there is also a unique length $\sqrt{\alpha'}$ that is invariant under T -duality which is a possible candidate for the *zero-point length*.

Let us consider a *closed* string winding w times around the compact dimension and freeze all the oscillator modes. By dropping the oscillator modes, we have reduced string dynamics to the motion of its center of mass. We will quantize the dynamics of the string center of mass (SCM), i.e. the Kaluza-Klein and winding modes. This is appropriate since we eventually want to obtain a corrected Feynman propagator in field theory. The latter can be intuitively thought of as giving the amplitude of propagation for a “particle”, the closest analogue of which is the string center of mass. We start from the propagation kernel of SCM in $4 + 1$ dimensions defined as

$$K[x - y; T] = \sum_{n=-\infty}^{\infty} \int_{z(0)=x}^{z(T)=y} \int_{x^5(0)=0}^{x^5(T)=n l_0} [Dz] [Dp] [Dx^5] [Dp_5] \times \exp \left[i \int_0^T d\tau \left(p_\mu \dot{x}^\mu + p_5 \dot{x}^5 - \frac{i}{2\mu_0} (p_\mu p^\mu + p_5 p^5) \right) \right] \quad (2)$$

where, $P_M \equiv (p_\mu, p_5)$ is the SCM five-momentum and μ_0 is a parameter introduced for dimensional reason which will not appear in the final result. In (2) we are summing over histories which in four dimensional space-time connect x and y in time T while, at the same time, they wind n times around x^5 .

In order to obtain the four dimensional propagator we need to integrate out the fifth-component of the paths.

This is most easily done by writing

$$\int_0^T p_5 \dot{x}^5 d\tau = [p_5 x^5]_0^T - \int_0^T d\tau x^5 \dot{p}_5 \quad (3)$$

and computing the path integral by

$$\int [\mathcal{D}x^5] \exp \left(-i \int_0^T d\tau x^5 \dot{p}_5 \right) = \delta[p_5] \quad (4)$$

This shows that p_5 is τ -independent, as it should be for a “free” particle. The final effect is to reduce functional integration over p_5 to an ordinary integral [5], but the compactification of the extra-dimension imposes a quantization condition on the SCM p_5 component of momentum, as: $p_5 = n \mu_0 (l_0/T)$ and we need to sum over the winding number n . This constraint can be implemented by introducing a Dirac-delta functional $\delta[p_5 - n \mu_0 (l_0/T)]$ in the path integral. Doing this and evaluating the integrals over the p_5 we get:

$$K(x - y; T) = 2 \sum_{n=1}^{\infty} \exp \left(\frac{in^2 \mu_0 l_0^2}{2T} \right) \times \int_{z(0)=x}^{z(T)=y} [Dz] [Dp] \exp \left(i \int_x^y p_\mu dz^\mu - i \int_0^T d\tau \frac{p^\mu p_\mu}{2\mu_0} \right) \quad (5)$$

In the summation over n , the $n = 0$ mode can be dropped since it does not sense the effect of the compact dimension which we are interested in. Further since only n^2 appears, the sum over negative integers is the same as that over positive integers. These features are already incorporated in the above result. The memory of the compact dimension is left over in the exponent involving summation over Kaluza-Klein modes.

The remaining integration over the four dimensional components can be done in a similar manner. Using

$$\int_{z(0)=x}^{z(T)=y} [Dz] \exp \left(i \int_x^y d(p_\mu z^\mu) - i \int_0^T d\tau z^\mu \dot{p}_\mu \right) = \delta[\dot{p}_\mu] \exp [i p_\mu z^\mu]_x^y \quad (6)$$

and integrating over p_μ , we get the final result:

$$K_{reg}(x - y; T) = -2 \left(\frac{\mu_0}{2i\pi T} \right)^2 \times \sum_{n=1}^{\infty} \exp \left\{ i \frac{\mu_0}{2T} \left[(x - y)^2 + n^2 l_0^2 \right] \right\} \quad (7)$$

The global normalization factor of the path integral measure is set in such a way to produce the correct form of the propagator in the limit $l_0 \rightarrow 0$.

The leading correction comes from the $n = 1$ which winds around the compact dimension just once. This reduces the expression in (7) to exactly the result obtained

in [3] from path integral duality. Thus the hypothesis introduced in [3], viz. the modification

$$\sum \exp[-l(x, y)] \rightarrow \sum \exp\left[-l + \frac{l_0^2}{l}\right] \quad (8)$$

captures the leading order corrections from string theory. The origin of zero-point length can be traced to the existence of compact extra-dimension. The ultraviolet divergences are, of course, cured because of the length scale l_0 .

Let us next compute $G(x - y)$ from the kernel by integrating over all possible values of the time lapse T with proper weight. In the standard Schwinger proper time approach to the propagator, this is done by using the weight $\exp(-im^2 T)$ where m is the physical mass of the particle which will emerge from the pole of the propagator. In our case, the SCM has a mass equal to the string tension times the total length of the string wrapped w times along x^5 , i.e. $M_{SCM} = w l_0 / \alpha'$ (In order to avoid any confusion, we remark that n counts the winding of the SCM histories summed in the path integral, while w refers to the string wrapping.) Thus, the Green function of SCM will be as

$$G_{reg}(x - y) \equiv \frac{i}{2\mu_0} \sum_w \int_0^\infty dT \exp\left(\frac{iT}{2\mu_0} m_0^2\right) \times \exp\left(\frac{iT}{2\mu_0} \frac{w^2 l_0^2}{\alpha'^2}\right) K_{reg}(x - y; T) \quad (9)$$

where m_0 is the mass of the particle in the limit $l_0 \rightarrow 0$, i.e. is the mass measured at low energies compared to threshold energy of the KK modes. The summation over w takes care of the KK modes and we have introduced m_0 to facilitate comparison with results of low energy field theory. (It can be set to zero for obtaining purely the SCM propagator). Inserting (7) into (9) we find

$$G_{reg}(x - y) = \frac{1}{(2\pi)^2} \sum_{w, n=1}^\infty \int_0^\infty \frac{du}{u^2} \exp\left(\frac{i u}{2} m_0^2\right) \times \exp\left(\frac{i u}{2} \frac{w^2 l_0^2}{\alpha'^2}\right) \exp\left\{\frac{i}{2u} \left[(x - y)^2 + n^2 l_0^2\right]\right\} \quad (10)$$

the unobservable parameter μ_0 has been eliminated by suitable rescaling $T \rightarrow u \equiv T/\mu_0$. The integral can be expressed in terms of the Bessel functions:

$$G_{reg}(x - y) = \frac{1}{(2\pi)^2} \sum_{w, n=1}^\infty \sqrt{\frac{m_0^2 + w^2 l_0^2 / \alpha'^2}{(x - y)^2 + n^2 l_0^2}} \times K_1\left(\sqrt{\left[(x - y)^2 + n^2 l_0^2\right] \left[m_0^2 + w^2 l_0^2 / \alpha'^2\right]}\right) \quad (11)$$

where $K_1(z)$ is a modified Bessel function. Inspection of (11) shows that $G_{reg}(x - y)$ is invariant under the transformation $(u/\alpha') \longleftrightarrow (\alpha'/u)$, $n \longleftrightarrow w$. This is the same as the T-duality (which makes the spectrum (1) invariant under $R \longleftrightarrow \alpha'/R$, $n \longleftrightarrow w$) with the identifications

$$(\alpha'/R^2) = (l_0^2/u); \quad (R^2/\alpha') = (u l_0^2/\alpha'^2) \quad (12)$$

which, in turn, requires precisely the identification $l_0^2 = \alpha'$. That is, the minimum length scale in string theory, set by $\sqrt{\alpha'}$, does get identified with the zero-point length, $l_0 \equiv \sqrt{\alpha'}$, when T-duality is properly encoded into the two-point Green function $G(x - y)$. This explains the T-duality origin of the zero-point length, in four dimensional spacetime.

The leading term in (11) is the one corresponding to $n = w = 1$. Writing $m^2 \equiv m_0^2 + l_0^2/\alpha'^2$, we get the leading terms to be

$$G_{reg}(x - y) \approx \frac{1}{(2\pi)^2} \left[\frac{m}{\sqrt{(x - y)^2 + n^2 l_0^2}} \times K_1\left(m \sqrt{(x - y)^2 + n^2 l_0^2}\right) \right] \quad (13)$$

The Fourier transform of this term gives the propagator in the momentum space, which is

$$G_{reg}(p^2) \approx \frac{1}{(2\pi)^2} \frac{l_0 K_1\left(l_0 \sqrt{p^2 + m^2}\right)}{\sqrt{p^2 + m^2}} \quad (14)$$

which is the result found earlier in [3].

Strictly speaking, the bosonic string can be consistently described only 26 dimensions. This requires the extension of 4 + 1 calculations to 4 + 22 dimensions. In a toroidal compactification scheme generalization of our results is easily achieved. There are compact dimensions of a priori different radii R_i and corresponding Kaluza-Klein and winding modes n_i and w_i . Nevertheless T-duality will relate all of them to the single parameter in the theory which is the string length $\sqrt{\alpha'}$. Thus, T-duality guarantees a unique zero-point length in the theory. Therefore, Green function can be obtained from (11) by the introduction of additional sum over compactified dimensions as:

$$n^2 \rightarrow \sum_{i=1}^{22} n_i^2; \quad w^2 \rightarrow \sum_{i=1}^{22} w_i^2 \quad (15)$$

We summarize our results as follows: Starting from the low energy end, we have the standard formulation of quantum field theory in which one can associate a propagator with a particle. Naively speaking, this propagator, $G(x - y)$ gives the amplitude for the particle to propagate from x to y . The ultra violet divergences of the standard

quantum field theory can be related to the fact that, G diverges as $x \rightarrow y$. One expects quantum gravitational effects to cure this behaviors and make G ultraviolet finite if there is a fundamental length scale in the theory. It was shown in [3] that such a result can be obtained if one postulates that the path integral amplitude used to compute G is invariant under the duality prescription $l \rightarrow L_P^2/l$. However, it was not clear how the duality and zero-point length of the spacetime (which leads to UV finiteness) are related. In this work, we have approached the problem from the string theory end, using the fact that standard formulation of string theory has both a fundamental length scale and T-duality. It is, however, not easy to obtain the low energy predictions of the string theory or the effective propagators for the low energy excitations, directly from the currently available formulation. This is because the low energy description of different particles is expected to emerge through a complex process in full fledged string theory. Though this makes an exact calculation difficult, one can capture the essential physics by treating the string center of mass as the analogue of the particle in the low energy limit and compute a propagator for it. We found that the winding over the compact dimensions leads to an infinite sum in this propagator. The $n = w = 1$ term gives the leading order contribution, which is identical to the result obtained in [3] using the prescription of path integral duality. This shows that, to the lowest order, the effects of string theory can be incorporated into ordinary quantum field theory by replacing $(x - y)^2$ by $(x - y)^2 + 4L_P^2$ in the Schwinger proper-time Kernel.

We incorporated the effect of low energy excitations by adding the corresponding phase term with a mass to the Schwinger proper-time integral. This allows us to define a low energy propagator and compare it with the results of earlier work. We stress that the incorporation

of the low mass excitations is logically separate from the calculation of propagator for SCM and the results for pure SCM propagator can be obtained by setting $m_0 = 0$.

The corrections to the standard path integral propagator obtained here [as well as in the earlier work] are non-perturbative by nature. That is, they cannot be obtained by a Taylor series expansion in L_P^2 . The expressions involved are like $[x^2 + L_P^2]^{-1}$ which is finite as $x^2 \rightarrow 0$; but if it is expanded in powers of L_P^2/x^2 , each term in the expansion will diverge as $x \rightarrow 0$.

One of the authors (T.P) thanks the Institute of Astronomy, Cambridge for hospitality where part of the work was carried out.

* Electronic address: e-mail: anais@ictp.trieste.it

† Electronic address: e-mail: spallucci@trieste.infn.it;
URL: <http://www-dft.ts.infn.it/~euro/>

‡ Electronic address: e-mail: nabhan@iucaa.ernet.in;
URL: <http://www.iucaa.ernet.in/~paddy>

- [1] L. J. Garay Int.J.Mod.Phys. **A10**, 145 (1995)
- [2] T. Padmanabhan Ann. Phys. (N.Y.) **165**, 38 (1985);
T. Padmanabhan Class. Quantum Grav. **4**, L107 (1987)
- [3] T. Padmanabhan Phys. Rev. Lett. **78**, 1854 (1997);
T. Padmanabhan, Phys. Rev. D, **57**, 6206 (1998).
- [4] M.B.Green, J.H.Schwartz, L.Brink Nucl. Phys. **B198**, 474 (1982)
K.Kikkawa, M.Yamasaki Phys. Lett. **B149**, 357 (1984)
N.Sakai,I.Senda Progr. Theor. Phys. **75**, 692 (1986)
V.P. Nair, A.Shapere, A.Strominger, F.Wilczek Nucl. Phys. **B287**, 402 (1987)
B.Sathiapalan Phys. Rev. Lett. **58**, 1597 (1987)
P.Ginsparg, C.Vafa Nucl. Phys. **B289**, 414 (1987)
A. Sen Int.J.Mod.Phys. **A9** 3707 (1994)
- [5] S.Ansoldi, A.Aurilia, E.Spallucci Eur.J.Phys. **21**, 1 (2000)